



# Gum disease creates pockets between the teeth and gums.

Gum disease can cause your gums to bleed or become infected, swollen, or tender. As a result, the infected gum starts to separate from the tooth.



The gap between the gum and the tooth is known as a pocket that can become deeper over time. And this may raise your risk of permanent damage.

## Where do your pockets fall on this chart?



### INDICATION

ARESTIN® is indicated as an adjunct to scaling and root planing (SRP) procedures for reduction of pocket depth in patients with adult periodontitis. ARESTIN® may be used as part of a periodontal maintenance program, which includes good oral hygiene and SRP.

### IMPORTANT SAFETY INFORMATION

ARESTIN® should not be used in any patient who has a known sensitivity to minocycline or tetracyclines. The use of drugs of the tetracycline class during tooth development may cause permanent discoloration of the teeth, and therefore should not be used in children or in pregnant or nursing women. Hypersensitivity reactions that included, but were not limited to, anaphylaxis, angioneurotic edema, urticaria, rash, swelling of the face, and pruritus have been reported with the use of ARESTIN®. In clinical trials, the most frequently reported nondental treatment-emergent adverse events were headache, infection, flu syndrome, and pain.

While ARESTIN® has been demonstrated along with SRP to reduce pocket depth and bacteria in patients with periodontal disease, it has not been proven in clinical trials to reduce tooth loss.

Please see full Prescribing Information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.